

Multiscale modeling of heterogeneous structures based on a localized model order reduction approach

P. Diercks*, A. Robens-Radermacher*, K. Veroy[†] and J.F. Unger*

* Bundesanstalt für Materialforschung und -prüfung (BAM)
Unter den Eichen 87
12205 Berlin

{philipp.diercks, annika.robens-radermacher, joerg.unger}@bam.de

[†] Centre for Analysis, Scientific Computing and Applications (CASA)
Department of Mathematics and Computer Science, TU Eindhoven
P.O. Box 513, 5600 MB Eindhoven, The Netherlands
k.p.veroy@tue.nl

Abstract: Many of today's problems in engineering demand reliable and accurate prediction of failure mechanisms of mechanical structures. Thus, it is necessary to take into account the heterogeneous structure on the smaller scale, to capture the underlying physical phenomena. However, this poses a great challenge to the numerical solution since the computational cost is significantly increased by resolving the smaller scale in the model. Moreover, in applications where scale separation as the basis of classical homogenization schemes does not hold, the influence of the smaller scale on the larger scale has to be modelled directly.

This work aims to develop an efficient concurrent methodology to model heterogeneous structures combining the variational multiscale method (VMM) [1] and model order reduction techniques (e.g. [2]). First, the influence of the smaller scale on the larger scale can be taken into account following the additive split of the displacement field as in the VMM. Here, also a decomposition of the global domain into subdomains, each containing a fine grid discretization of the smaller scale, is introduced. Second, local reduced approximation spaces for the smaller scale solution are constructed by exploring possible solutions for each subdomain based on the concept of oversampling [3]. The associated transfer operator is approximated by random sampling [4]. Herein, we propose to incorporate the actual physical behaviour of the structure of interest in the training data by drawing random samples from a multivariate normal distribution with the solution of a reduced global problem as mean. The local reduced spaces are designed such that local contributions of each subdomain can be coupled in a conforming way. Thus, the resulting global system is sparse and reduced in size compared to the direct numerical simulation, leading to a faster solution of the problem.

The authors gratefully acknowledge financial support by the German Research Foundation (DFG), project number 394350870, and by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (ERC Grant agreement No. 818473).

Keywords: Multiscale Method, Variational Multiscale Method, Domain Decomposition, Model Order Reduction

References

- [1] Thomas J.R. Hughes, Gonzalo R. Feijóo, Luca Mazzei, and Jean-Baptiste Quinicy. The variational multiscale method—a paradigm for computational mechanics. *Comput. Method. Appl. M.*, 166(1-2):3–24, November 1998.
- [2] Jan S Hesthaven, Gianluigi Rozza, and Benjamin Stamm. *Certified Reduced Basis Methods for Parametrized Partial Differential Equations*. SpringerBriefs in Mathematics. Springer International Publishing, 2016.
- [3] Thomas Y. Hou and Xiao-Hui Wu. A multiscale finite element method for elliptic problems in composite materials and porous media. *J. Comput. Phys.*, 134(1):169–189, June 1997.
- [4] Andreas Buhr and Kathrin Smetana. Randomized local model order reduction. *SIAM J. Sci. Comput.*, 40(4):A2120–A2151, January 2018.